Day	Treatment	
	"Single" (two-injection) protocol	"Split" (three-injection) protocol
Day 0	Dog diagnosed and verified as heartworm positive: • Positive antigen (Ag) test verified with microfilaria (MF) test • If no microfilariae are detected, confirm with 2nd Ag test from a different manufacturer Begin exercise restriction. The more pronounced the signs, the stricter the exercise restriction If the dog is symptomatic: • Stabilize with appropriate therapy and nursing care • Prednisone prescribed at 0.5 mg/kg BID 1st week, 0.5 mg/kg SID 2nd week, 0.5 mg/kg EOD 3rd and 4th weeks	

Da 1	ау	 Administer heartworm preventive. If microfilariae are detected, pretreat with antihistamine and glucocorticosteroid, if not already on prednisone, to reduce risk of anaphylaxis Observe for at least 8 hours for signs of reaction 	
	ays 28	Administer doxycycline 10 mg/kg BID for 4 weeks. • Reduces pathology associated with dead heartworms • Disrupts heartworm transmission	
Da 30	-	Administer heartworm preventative	
Da 60	•	Administer heartworm preventive. First melarsomine injection 2.5 mg/kg intramuscularly (IM) Prescribe prednisone 0.5 mg/kg BID 1st week, 0.5 mg/kg SID 2nd week, 0.5 mg/kg EOD 3rd and 4th weeks. Decrease activity level even further.	Administer heartworm preventive. First melarsomine injection 2.5 mg/kg intramuscularly (IM) Prescribe prednisone 0.5 mg/kg BID 1st week, 0.5 mg/kg SID 2nd week, 0.5 mg/kg EOD 3rd and 4th weeks. Decrease activity level even further.

	 Cage restriction/on leash when using yard 	 Cage restriction/on leash when using yard
Day 61	Second melarosamine injection 2.5mg/kg IM	
Day 90	Administer heartworm preventive.	Administer heartworm preventative. Second melarsomine injection 2.5 mg/kg IM
Day 91	Administer heartworm preventative.	Third melarsomine injection 2.5 mg/kg IM Prescribe prednisone 0.5 mg/kg BID 1st week, 0.5 mg/kg SID 2nd week, 0.5 mg/kg EOD 3rd and 4th weeks. Continue exercise restriction for 6 to 8 weeks following last melarsomine injections.
Day 120	Establish year-round heartworm prevention.	
Day ~330	Antigen test 9 months after completion; screen for microfilariae.	

Day	Antigen test 9 months after
~365	completion; screen for microfilariae.